



## **A molecular method to differentiate between three genotypes of trichuris recovered from humans and pigs**

Al-Jubury, Azmi; Nissen, Sofie; Olsen, Annette; Christensen, Henrik; Thamsborg, Stig Milan; Nejsun, Peter

*Publication date:*  
2010

*Document version*  
Publisher's PDF, also known as Version of record

*Citation for published version (APA):*  
Al-Jubury, A., Nissen, S., Olsen, A., Christensen, H., Thamsborg, S. M., & Nejsun, P. (2010). *A molecular method to differentiate between three genotypes of trichuris recovered from humans and pigs*. Abstract from Joint Spring Symposium 2010, Danish Society for Parasitology and Danish Society for Tropical Medicine & International Health, Denmark.

## JOINT SPRING SYMPOSIUM 2010

Danish Society for Parasitology and  
Danish Society for Tropical Medicine & International Health

### A MOLECULAR METHOD TO DIFFERENTIATE BETWEEN THREE GENOTYPES OF *TRICHURIS* RECOVERED FROM HUMANS AND PIGS

Azmi Al-Jubury<sup>1</sup>, Sofie Nissen<sup>1</sup>, Annette Olsen<sup>1</sup>, Henrik Christensen<sup>2</sup>, Stig M. Thamsborg<sup>1</sup> and Peter Nejsum<sup>1</sup>

<sup>1</sup> Parasitology, health and development; <sup>2</sup> Microbiology, Department of Veterinary Disease Biology, Faculty of Life Science, University of Copenhagen.

#### Abstract

The whipworms *Trichuris trichiura* and *T. suis* in humans and pigs, respectively, are believed to be closely related but different species. Despite several attempts no reliable morphological characters have been identified which can discriminate worms from the two hosts. Recent sequence analysis has suggested that 3 major ITS-2 genotypes are present in *Trichuris* obtained from human and pigs; namely a type 1 and 2 and a heterozygote. We have used this information to develop a PCR-linked Restriction Fragment Length Polymorphism (PCR-RFLP) that can identify the three genotypes. The method was applied on worms obtained from pigs collected in Uganda, USA, Tanzania, Jamaica and Denmark and worms obtained from humans collected in Uganda, Jamaica and China. The distribution of the three genotypes (type 1, type 2 and heterozygote) was 2, 0 and 57, respectively, in 59 worms recovered from pigs and 26, 2 and 1, respectively in 29 worms recovered from human. The results clearly suggest that *Trichuris* in humans and pigs belong to two different populations ( $\chi^2 = 49.5$ ;  $P < 0.001$ ). However, we have also found evidence for a few cases of cross-infections and the existence of heterozygote worms, the latter suggesting cases of hybridization or retention of ancestral polymorphism.